## CBSE TEST PAPER-01 CLASS - XI BIOLOGY (Plant Kingdom)

### **General Instruction:**

- All questions are compulsory.
- Question No. 1 to 3 carry one marks each. Question No. 4 to 6 carry two marks each. Question No. 7 and 8 carry three marks each. Question No. 9 carry five marks.
- 1. Name the group of seedless vascular plants.
- 2. Which pigment is responsible for red colour of red algae?
- 3. Define a cone?
- 4. Comment on the features that led to the dominance of vascular plants?
- 5. Distinguish between Red algae & brown algae?
- 6. Enlist few salient features of dicot plants?
- 7. Briefly explain the structure of prothallus of a fern?
- 8. Point out differences between the mode of sexual reproduction of moss & fern?
- 9. Discuss the various life cycles of a green algae?

# CBSE TEST PAPER-01 CLASS - XI BIOLOGY (Plant Kingdom) [ANSWERS]

1. Pteridophytes.

2. Phycobilin, phycoerythrin & phycocyanin.

3. Cones are the organs of Reproduction in Gymnosperms. They are also called the fruiting body which consists of micro & megasporophyll. Microsporohylls are the structures of Male sex organ while megasporophylls are the structures of female sex organ.

4. Three important features have to dominance of Angiosperm:-

i) Development of well established deep roots capable of penetrating the soil to absorb waterii) Development of water- proofing material eg. cutin on aerial surfaces, to reduce water loss through evaporation.

iii) Development of strong woody material as anchor & support above ground structures.

5.

RED ALGAE	BROWN ALGAE
i) Mostly unicallular & microscopic	i) filamentous & heterotrichous and
i) Mostry unicendial & incroscopic	macroscopic also
) Phycoerythrin, phycocyanin & phycobilin	
pigments are present.	n) iucoxantinii pignieni is present.
li jij) Posorvo food matorial is Floridian starch	iii) Reserve food material is Laminarian
II III) Reselve 1000 material is i loridian starch	starch.
iv) chlorophyll 'a' present	iv) chlorophyll 'a' absent
v) eg. Gelidium, Polysiphonia	v) Laminaria, Fucus & Sargassum

6. The dicotyledons are characterized by either woody or herbaceous habit, They have well developed tap root system. Their floral parts usually are in Tetramerous or Pentamerous. their leaves have Reticulate venation. In the stem the vascular bundles are Open, Collateral and arranged in a circle or ring. Vascular bundles are wedge shaped and possess Vascular cambium. The dicots have two cotyledons in their seeds.

### 7. Prothallus of Fern:-



- i) It is a heart-shaped structure, autotrophic and independant
- ii) The sex organs are present on the lower surface of the prothallus below the apical notch.
- iii) Male Sex organs are antheridia & the female ones are archaegonia.
- iv) Prothallus is produced from the meiospore.
- v) Below the sex organs are rhizoids
- vi) Archegonia are flask shaped but antheridia are globose.
- vii) Male & female gametes are produced in antheridia & archegonia respectively

#### 8.

MOSS (Bryophyte)	Fern (pteridophytes)	
i) Sex organs are borne on	i) Sex organs are borne on an inconspicuous gametophyte or	
the Primary gametophytic	prothallus which represents an alternate and secondary phase	
plant body.	to sporophytic phase which is the primary plant body.	
ii). Antheridia are well		
developed & often possess	ii) Anthonidia and loss devialanced & mostly deviaid of a stall	
a stalk caled	n). Antheriula are less developed & mostly devold of a stalk.	
Antheridiophore		
iii). Antheridial jacket		
made up of a layer of	iii). Antheridial jacket is mostly made up of only 3 – cells.	
several cells.		
iv). Sperms biflagellate	iv). Multiflagellate sperms	
v). Archegonia often have		
stalk caled	v). Archegonia do not have stalk	

archegoinophore	
vi). Neck is 6 - rowed	vi). Neck is 4 – rowed.

9. There are three types of life cycle are found in green algae:-

a) HAPLONTIC LIFE CYCLE:- The dominant phase is haploid. Diploid state is found only in the form of zygote. Meiosis takes place at time of its germination. It has Zygotic meiosis Eg. *Ulothrix, Spirogyra*.

b) DIPLONTIC LIFE CYCLE:- The dominant phase of the alga is diploid. It gives rise to haploid gametes through meiosis leading to gametic meiosis Gametes unit & the zygote regenerates diploid phase. Gametes alone are haploid Eg. Diatoms

c) DIPLOHAPLONTIC LIFE CYCLE:- It has well developed multicellular haploid & diploid phase. These are respectively called gametophyte & sporophyte. Haploid gametophyte produce haploid gametes. Fusion product of gametes grows directly into diploid sporophytes. Sporophytes produce haploid spores by meiosis. The meiospores germinate into new gametophyte. Eg. *Ulva, Cladophora* 

